

Polypeptide

Field of the Invention

- 5 The present invention relates to a tumour-associated antigen (TAA) useful for eliciting an anti-tumour immunotherapeutic response in subjects. In particular, the invention relates to 5T4 antigen and its use in immunotherapy.

Background to the Invention

10

- A number of oncofoetal or tumour-associated antigens (TAAs) have been identified and characterised in human and animal tumours. In general, TAAs are antigens expressed during foetal development which are downregulated in adult cells, and are thus normally absent or present only at very low levels in adults. Tumour cells have
15 been observed to resume expression of TAAs, and the application of TAAs for tumour diagnosis, targeting and immunotherapy has therefore been suggested.

- In particular, the recent cloning of tumour antigens recognised by T cells has caused considerable interest in the development of antigen specific cancer vaccines. However,
20 many tumour associated antigens are non-mutated, poorly immunogenic tissue differentiation antigens. Their weak immunogenicity may be due to self tolerance. Thus they are rarely indicated as antigenic peptides suitable for raising an immune response.

- 25 Notwithstanding this, some tumour associated antigens are found to be regularly associated with tumours in a large number of individuals. Such antigens are especially attractive candidates for use in vaccines. They include the melanoma differentiation antigens (MDA), melanoma antigens which are recognised by T lymphocytes as well as several proteins in the MAGE family. However, as indicated by results from clinical
30 trials obtained to date, inducing therapeutic T cells to these antigens has proved extremely difficult. One reason for the apparent hyporesponsiveness of the human immune system to many tumour antigens may be that they are normal, non-mutated

self antigens, expressed on normal tissues as well as on tumour cells. The immune system is not able to differentiate the tumour antigen on a tumour cell from ordinary, self proteins.

- 5 A major barrier to the application of tumour immunotherapy approaches using non-mutated self cellular antigens is thus apparently the breaking of tolerance to such an antigen. For example, a murine zona pellucida antigen expressed by a murine poxvirus recombinant was able to induce infertility in mice. These data indicate that though the breaking of tolerance using recombinant pox viruses expressing self
10 antigens is possible, there is still a requirement to optimise their efficacy such that the active treatment of established tumours becomes possible.

The TAA 5T4 (see WO 89/07947) has been extensively characterised. It is a 72kDa glycoprotein expressed widely in carcinomas, but having a highly restricted expression
15 pattern in normal adult tissues (see Table 1). It appears to be strongly correlated to metastasis in colorectal and gastric cancer. The full nucleic acid sequence of human 5T4 is known (Myers *et al.*, 1994 J Biol Chem 169: 9319-24).

Table 1

Distribution of Human 5T4		
Tumour	5T4	Frequency
Type	(%)	
Breast	84	
Ovarian	71	
Gastric	74	
Colorectal	85	

20

(Starzynska *et al.*, Eur J Gastroenterol Hepatol 1998 Jun;10(6):479-84; Starzynska *et al.*, Br J Cancer 1994 May;69(5):899-902; Starzynska *et al.*, Br J Cancer 1992 Nov;66(5):867-9)

Although 5T4 has been proposed as a marker, with possible mechanistic involvement, for tumour progression and metastasis potential (Carsberg *et al.*, (1996) Int J Cancer 1996 Sep 27;68(1):84-92), 5T4 has not been proposed for use as an immunotherapeutic agent. The breaking of immune tolerance to 5T4, which is itself
 5 expressed in a restricted manner in adult tissues, has not been demonstrated. Thus, it could not be predicted whether 5T4 could prove to be an effective antigen for immunotherapy against cancer.

10 Summary of the Invention

If a successful therapeutic outcome is to be achieved, an immunotherapeutic approach to cancer treatment depends on a number of factors. These include the ability to elicit a cytotoxic T-lymphocyte (CTL) response, the ability to elicit an antibody response
 15 and, importantly, the ability to break immune tolerance in a subject. It has now been demonstrated that immunisation of subjects with 5T4 results in a successful immunotherapeutic response as judged by the above. In particular, immunisation with 5T4 has been shown to elicit an antibody response.

20 Accordingly, the present invention provides a viral vector expressing a nucleic acid encoding 5T4 antigen.

Expression of 5T4 antigen in a subject is effective in eliciting an immunotherapeutic anti-tumour response. Preferably, the viral vector favours CTL responses to expressed
 25 antigens, and is advantageously a poxvirus vector, such as a vaccinia virus vector. Further vectors, both viral and non-viral, which are suitable for delivering 5T4 antigen are described below.

As used herein, a "vector" may be any agent capable of delivering or maintaining
 30 nucleic acid in a host cell, and includes viral vectors, plasmids, naked nucleic acids, nucleic acids complexed with polypeptide or other molecules and nucleic acids immobilised onto solid phase particles. Such vectors are described in detail below. It

will be understood that the present invention, in its broadest form, is not limited to any specific vector for delivery of the 5T4-encoding nucleic acid.

A "nucleic acid", as referred to herein, may be DNA or RNA, naturally-occurring or synthetic, or any combination thereof. Nucleic acids according to the invention are limited only in that they serve the function of encoding 5T4 antigen in such a way that it may be translated by the machinery of the cells of a host organism. Thus, natural nucleic acids may be modified, for example to increase the stability thereof. DNA and/or RNA, but especially RNA, may be modified in order to improve nuclease resistance of the members. For example, known modifications for ribonucleotides include 2'-O-methyl, 2'-fluoro, 2'-NH₂, and 2'-O-allyl. The modified nucleic acids according to the invention may comprise chemical modifications which have been made in order to increase the *in vivo* stability of the nucleic acid, enhance or mediate the delivery thereof, or reduce the clearance rate from the body. Examples of such modifications include chemical substitutions at the ribose and/or phosphate and/or base positions of a given RNA sequence. See, for example, WO 92/03568; U.S. 5,118,672; Hobbs *et al.*, (1973) *Biochemistry* 12:5138; Guschlbauer *et al.*, (1977) *Nucleic Acids Res.* 4:1933; Schibaharu *et al.*, (1987) *Nucleic Acids Res.* 15:4403; Pieken *et al.*, (1991) *Science* 253:314, each of which is specifically incorporated herein by reference.

5T4 antigen is "expressed" in accordance with the present invention by being produced in the cells of a host organism as a result of translation, and optionally transcription, of the nucleic acid encoding 5T4. Thus, 5T4 is produced *in situ* in the cell. Since 5T4 is a transmembrane protein, the extracellular portion thereof is displayed on the surface of the cell in which it is produced. If necessary, therefore, the term "expression" includes the provision of the necessary signals to ensure correct processing of 5T4 such that it is displayed on the cell surface and can interact with the host immune system.

30

As used herein, the term "polypeptide" refers to a polymer in which the monomers are amino acids and are joined together through peptide or disulphide bonds.

“Polypeptide” refers to a full-length naturally-occurring amino acid chain or a fragment thereof, such as a selected region of the polypeptide that is of interest in a binding interaction, or a synthetic amino acid chain, or a combination thereof. “Fragment thereof” thus refers to an amino acid sequence that is a portion of a full-length polypeptide, between about 8 and about 500 amino acids in length, preferably about 8 to about 300, more preferably about 8 to about 200 amino acids, and even more preferably about 10 to about 50 or 100 amino acids in length. Additionally, amino acids other than naturally-occurring amino acids, for example β -alanine, phenyl glycine and homoarginine, may be included. Commonly-encountered amino acids which are not gene-encoded may also be used in the present invention.

5T4 antigen is the polypeptide known as 5T4 and characterised, for example, in WO89/07947. In a preferred aspect, 5T4 is human 5T4 as characterised by Myers *et al* *ibid.*, the sequence of which appears in GenBank at accession no. Z29083 and is set out herein as SEQ. ID. No. 1. The invention however comprises species and allelic variations of 5T4, including canine 5T4 set forth herein at SEQ. ID. No. 3 and mouse 5T4 set forth herein at SEQ. ID. No. 2 (GenBank Accession no. AJ012160), as well as fragments, preferably distinct epitopes, and variants thereof comprising amino acid insertions, deletions or substitutions which retain the antigenicity of 5T4. Such fragments and variants are described in greater detail below.

In a second aspect, the present invention relates to a modified 5T4 antigen. A “modified” antigen, as used herein, is a 5T4 polypeptide which has been truncated, extended or otherwise mutated such that it differs from naturally-occurring 5T4. It has been found that peptide fragments derived from 5T4 are able to function as 5T4-specific antigenic determinants. Such peptides are able to bind HLA molecules and to induce CTL responses against wild-type 5T4 in subjects, often more effectively than full-length 5T4. Moreover, 5T4 peptides may be mutated, by amino acid insertion, deletion or substitution; mutated peptides advantageously bind even more effectively to HLA and elicit an even more potent CTL response in subjects. Peptides may be any length, but are advantageously between 5 and 25 amino acids, preferably between 6 and 15 amino acids, and advantageously about 9 amino acids in length.

Modified peptides are advantageously HLA CTL epitopes of 5T4. Modification of such epitopes may be performed based on predictions for more efficient CTL induction derived using the program "Peptide Binding Predictions" devised by K. Parker (NIH) which may be found at http://www-bimas.dcrn.nih.gov/cgi-bin/molbio/ken_parker_comboform (see also Parker, K. C *et al.* 1994.J.Immunol. 152:163).

In a preferred aspect, a "modified" 5T4 peptide includes peptides which have been bound or otherwise associated to transporter peptides or adjuvants, in order to increase their ability to elicit an immune response. For example, peptides may be fused to TAP independent transporter peptides for efficient transport to HLA and interaction with HLA molecules to enhance CTL epitopes (for review see Yewdell *et al.*, 1998 J Immunother 21:127-31; Fu *et al.*, (1998) J Virol 72:1469-81).

In a third aspect, the present invention provides a method for eliciting an immune response in a subject, comprising the steps of immunising the subject with a nucleic acid encoding 5T4 antigen, and expressing the 5T4 antigen in the subject.

An immune response is elicited, as stated, by immunisation with 5T4-expressing nucleic acid. Immunisation may be elicited through the administration of a "priming" agent comprising an antigen followed by a secondary or "boosting" agent comprising additional antigen which is administered to the immune system after it has been efficiently primed with the priming agent.

The vector employed for immunisation may be any vector, viral or non-viral. The 5T4 antigen used, whether full length 5T4 or peptides thereof, may be modified and may be homologous (i.e. derived from the same species as the subject) or heterologous in origin.

Preferably, the immune response elicited is a CTL response which involved the activation of cytotoxic T-lymphocytes which are 5T4 specific.

Advantageously, the response is an anti-tumour immunotherapeutic response which is effective to inhibit, arrest or reverse the development of a tumour in a subject.

- 5 In a fourth aspect, the present invention provides the use of a 5T4 antigen in the preparation of a composition for the immunotherapy of a tumour in a subject.

Advantageously, immunisation with a 5T4 antigen is capable of breaking immune tolerance to 5T4 in a subject.

10

According to a fifth aspect, the present invention provides a vaccine composition comprising 5T4 antigen. The vaccine composition may comprise a homologous 5T4 antigen, a heterologous 5T4 antigen or a mutant 5T4 antigen.

- 15 5T4 antigen-containing vaccines are useful for immunisation against, or therapy of, tumours, in a manner analogous to the use of 5T4-encoding nucleic acids for the same purposes.

Advantageously, the vaccine composition comprises one or more adjuvants.

20

In a sixth aspect, the invention comprises an expression vector encoding a 5T4 antigen, which vector is useful for the expression of 5T4 and the production of 5T4 antigen suitable for use in a vaccine composition. The vector may be a prokaryotic or eukaryotic vector, and is advantageously a vector capable of expressing 5T4 in
25 mammalian cells.

The 5T4 antigen may be from any source, and may be a modified 5T4 antigen, for example as set forth herein.

- 30 In a seventh aspect, the present invention provides the use of a 5T4 antigen in the preparation of a composition for immunising a subject. Immunisation using a 5T4

comprises administering to the subject an immunologically effective amount of the vaccine composition according to the fifth aspect of the invention.

5 In an eighth aspect, the present invention provides the use of a 5T4 antigen in the preparation of a composition for the sterilisation of a subject. The administration of 5T4 antigen may be effective in causing sterilisation of subjects. Preferably, the subject is a female subject.

10 The invention further relates to the use of 5T4 targeting molecules, such as anti-5T4 antibodies, for example anti-5T4 scFvs. These antibodies may be used to (i) to target natural or exogenous 5T4 *in situ* and/or (ii) deliver immune enhancer molecules, such as B7.1, to natural or exogenous 5T4 *in situ* (Carroll *et al.* (1998) J Natl Cancer Inst 90(24):1881-7). This potentiates the immunogenicity of 5T4 in the subject. The present invention also relates to the sequential use of a vector encoding a 5T4 antigen and anti-5T4 antibodies, for example an anti-5T4 scFvs. The anti-5T4 scFvs antibodies may be administered as naked DNA encoding the antibodies (for example, in a plasmid comprising the encoding DNA together with a short promoter region to control its production), in an expression vector (which may be viral or non-viral) comprising the encoding sequence or in a protein form. Thus, the invention provides a vector encoding a 5T4 antigen and an agent capable of binding 5T4 which is optionally fused with an immunostimulatory molecule, for separate, such as sequential use, in the treatment of tumours.

25 In a further embodiment, the invention encompasses a combination therapy including enzyme/prodrug therapy and immunotherapy with 5T4. For example, the enzyme/prodrug therapy may comprise intratumoural or systemic delivery of P450, delivered optionally using an retroviral or lentiviral vector, and cyclophosphamide (CPA) followed by systemic immunotherapeutic induction with 5T4.

30 Thus, the invention further relates to a vector encoding 5T4 antigen a prodrug/enzyme combination, for separate, simultaneous separate or combined use in the treatment of tumours.

In a further embodiment, 5T4 or 5T4 peptides may be fused to hepatitis B core antigen to enhance T helper and antibody responses (Schodel *et al.*, 1996 Intervirology 39:104-10).

5 In accordance with a ninth aspect of the invention, therefore, there is provided a recombinant poxvirus vector from which at least one immune evasion gene has been deleted, which comprises a nucleic acid sequence encoding a tumour-associated antigen (TAA).

10 TAAs are weakly immunogenic, being recognised as "self" by the immune system and thus tolerated to a large extent. Although the use of poxvirus vectors is able to cause the antigens to be presented such that this tolerance may be overcome at least in part, the immunogenic effect observed with most poxvirus vectors is limited. It is thought that the deletion of immune evasion genes, naturally present in poxviruses, may have a
15 beneficial effect in vaccination with TAAs. Poxvirus vectors having deleted immune evasion genes may be capable of breaking immune tolerance to encoded self antigens, including TAAs, thus enabling a host to raise an immune response to a poorly immunogenic or other self antigen.

20 In a tenth aspect, the present invention provides a method for eliciting an immune response in a mammal, comprising administering to the mammal a recombinant poxvirus vector according to the ninth aspect of the invention, thereby eliciting an immune response to the TAA in the mammal.

25 Antigens such as TAAs are known to rely on the generation of a CTL response in order to provide a protective or therapeutic effect in a subject, which is dependent on processing of antigen via the MHCI pathway. Long-term antigen expression is thought to lead to increased longevity of high level CTL. In an eleventh aspect of the invention, there is provided a poxvirus having a reduced lytic activity for the
30 enhancement of a CTL response to an antigen in a subject.

In a twelfth aspect, the present invention provides the use of a recombinant poxvirus vector according to the ninth or eleventh aspect of the invention, to elicit an immune response in a mammal against a TAA.

- 5 In a thirteenth aspect, the invention provides a 5T4 antigen for use as a tumour-associated target in immunotherapy.

In a fourteenth aspect, the invention provides the use of a recombinant poxvirus vector from which at least one immune evasion gene has been deleted or mutated,
10 which comprises a nucleic acid sequence encoding a weak immunogen, to break immune tolerance in a mammal against the weak immunogen and elicit an immune response thereto.

Other aspects of the present invention are presented in the accompanying claims and in
15 the following description and discussion. These aspects are presented under separate section headings. However, it is to be understood that the teachings under each section heading are not necessarily limited to that particular section heading.

Detailed Description of the Invention

20

Vectors for delivery or expression of 5T4 antigen

5T4 polypeptides in accordance with the present invention can be delivered by viral or non-viral techniques.

25

Non-viral delivery systems include but are not limited to DNA transfection methods. Here, transfection includes a process using a non-viral vector to deliver a 5T4 gene to a target mammalian cell.

30 Typical transfection methods include electroporation, nucleic acid biolistics, lipid-mediated transfection, compacted nucleic acid-mediated transfection, liposomes, immunoliposomes, lipofectin, cationic agent-mediated, cationic facial amphiphiles

(CFAs) (Nature Biotechnology 1996 14; 556), multivalent cations such as spermine, cationic lipids or polylysine, 1, 2,-bis (oleoyloxy)-3-(trimethylammonio) propane (DOTAP)-cholesterol complexes (Wolff and Trubetskoy 1998 Nature Biotechnology 16: 421) and combinations thereof.

- 5 Viral delivery systems include but are not limited to adenovirus vectors, adeno-associated viral (AAV) vectors, herpes viral vectors, retroviral vectors, lentiviral vectors or baculoviral vectors, venezuelan equine encephalitis virus (VEE), poxviruses such as: canarypox virus (Taylor *et al* 1995 Vaccine 13:539-549), entomopox virus (Li Y *et al* 1998 XIIth International Poxvirus Symposium p144. Abstract), penguin pox
10 (Standard *et al.* J Gen Virol. 1998 79:1637-46) alphavirus, and alphavirus based DNA vectors.

- Examples of retroviruses include but are not limited to: murine leukaemia virus (MLV), human immunodeficiency virus (HIV), equine infectious anaemia virus
15 (EIAV), mouse mammary tumour virus (MMTV), Rous sarcoma virus (RSV), Fujinami sarcoma virus (FuSV), Moloney murine leukaemia virus (Mo-MLV), FBR murine osteosarcoma virus (FBR MSV), Moloney murine sarcoma virus (Mo-MSV), Abelson murine leukaemia virus (A-MLV), Avian myelocytomatosis virus-29 (MC29), and Avian erythroblastosis virus (AEV).

- 20 A detailed list of retroviruses may be found in Coffin *et al* ("Retroviruses" 1997 Cold Spring Harbour Laboratory Press Eds: JM Coffin, SM Hughes, HE Varmus pp 758-763).

- 25 Lentiviruses can be divided into primate and non-primate groups. Examples of primate lentiviruses include but are not limited to: the human immunodeficiency virus (HIV), the causative agent of human auto-immunodeficiency syndrome (AIDS), and the simian immunodeficiency virus (SIV). The non-primate lentiviral group includes the prototype "slow virus" visna/maedi virus (VMV), as well as the related caprine
30 arthritis-encephalitis virus (CAEV), equine infectious anaemia virus (EIAV) and the more recently described feline immunodeficiency virus (FIV) and bovine immunodeficiency virus (BIV).

A distinction between the lentivirus family and other types of retroviruses is that lentiviruses have the capability to infect both dividing and non-dividing cells (Lewis *et al* 1992 EMBO. J 11: 3053-3058; Lewis and Emerman 1994 J. Virol. 68: 510-516). In contrast, other retroviruses - such as MLV - are unable to infect non-dividing cells such as those that make up, for example, muscle, brain, lung and liver tissue.

The vector of the present invention may be configured as a split-intron vector. A split intron vector is described in PCT patent applications WO 99/15683 and WO 99/15684.

10

If the features of adenoviruses are combined with the genetic stability of retroviruses/lentiviruses then essentially the adenovirus can be used to transduce target cells to become transient retroviral producer cells that could stably infect neighbouring cells. Such retroviral producer cells engineered to express 5T4 antigen can be implanted in organisms such as animals or humans for use in the treatment of angiogenesis and/or cancer.

15

Poxvirus Vectors

Poxvirus vectors are preferred for use in the present invention. Pox viruses are engineered for recombinant gene expression and for the use as recombinant live vaccines. This entails the use of recombinant techniques to introduce nucleic acids encoding foreign antigens into the genome of the pox virus. If the nucleic acid is integrated at a site in the viral DNA which is non-essential for the life cycle of the virus, it is possible for the newly produced recombinant pox virus to be infectious, that is to say to infect foreign cells and thus to express the integrated DNA sequence. The recombinant pox virus prepared in this way can be used as live vaccines for the prophylaxis and/or treatment of pathologic and infectious disease.

25

Expression of 5T4 in recombinant pox viruses, such as vaccinia viruses, requires the ligation of vaccinia promoters to the nucleic acid encoding 5T4. Plasmid vectors (also called insertion vectors), have been constructed to insert nucleic acids into vaccinia

30

virus through homologous recombination between the viral sequences flanking the nucleic acid in a donor plasmid and homologous sequence present in the parental virus (Mackett *et al* 1982 PNAS 79: 7415-7419). One type of insertion vector is composed of: (a) a vaccinia virus promoter including the transcriptional initiation site; (b) several
 5 unique restriction endonuclease cloning sites located downstream from the transcriptional start site for insertion of nucleic acid; (c) nonessential vaccinia virus sequences (such as the Thymidine Kinase (*TK*) gene) flanking the promoter and cloning sites which direct insertion of the nucleic acid into the homologous nonessential region of the virus genome; and (d) a bacterial origin of replication and
 10 antibiotic resistance marker for replication and selection in *E. Coli*. Examples of such vectors are described by Mackett (Mackett *et al* 1984, J. Virol. 49: 857-864).

The isolated plasmid containing the nucleic acid to be inserted is transfected into a cell culture, e.g., chick embryo fibroblasts, along with the parental virus, e.g., poxvirus.
 15 Recombination between homologous pox DNA in the plasmid and the viral genome respectively results in a recombinant poxvirus modified by the presence of the promoter-gene construct in its genome, at a site which does not affect virus viability.

As noted above, the nucleic acid is inserted into a region (insertion region) in the virus
 20 which does not affect virus viability of the resultant recombinant virus. Such regions can be readily identified in a virus by, for example, randomly testing segments of virus DNA for regions that allow recombinant formation without seriously affecting virus viability of the recombinant. One region that can readily be used and is present in many viruses is the thymidine kinase (*TK*) gene. For example, the *TK* gene has been
 25 found in all pox virus genomes examined [leporipoxvirus: Upton, *et al* J. Virology 60:920 (1986) (Shope fibroma virus); capripoxvirus: Gershon, *et al* J. Gen. Virol. 70:525 (1989) (Kenya sheep-1); orthopoxvirus: Weir, *et al* J. Virol 46:530 (1983) (vaccinia); Esposito, *et al* Virology 135:561 (1984) (monkeypox and variola virus); Hruby, *et al* PNAS, 80:3411 (1983) (vaccinia); Kilpatrick, *et al* Virology 143:399
 30 (1985) (Yaba monkey tumour virus); avipoxvirus: Binns, *et al* J. Gen. Virol 69:1275 (1988) (fowlpox); Boyie, *et al* Virology 156:355 (1987) (fowlpox); Schnitzlein, *et al*

J. Virological Method, 20:341 (1988) (fowlpox, quailpox); entomopox (Lytvyn, *et al* J. Gen. Virol 73:3235-3240 (1992)].

5 In vaccinia, in addition to the TK region, other insertion regions include, for example, HindIII M.

In fowlpox, in addition to the TK region, other insertion regions include, for example, BamHI J [Jenkins, *et al* AIDS Research and Human Retroviruses 7:991-998 (1991)] the *EcoRI-HindIII* fragment, *BamHI* fragment, *EcoRV-HindIII* fragment, *BamHI* fragment and the *HindIII* fragment set forth in EPO Application No. 0 308 220 A1.
10 [Calvert, *et al* J. of Virol 67:3069-3076 (1993); Taylor, *et al* Vaccine 6:497-503 (1988); Spehner, *et al* (1990) and Bournsnel, *et al* J. of Gen. Virol 71:621-628 (1990)].

In swinepox preferred insertion sites include the thymidine kinase gene region.

15 A promoter can readily be selected depending on the host and the target cell type. For example in poxviruses, pox viral promoters should be used, such as the vaccinia 7.5K, or 40K or fowlpox C1. Artificial constructs containing appropriate pox sequences can also be used. Enhancer elements can also be used in combination to increase the level of expression. Furthermore, the use of inducible promoters, which are also well
20 known in the art, are preferred in some embodiments.

Foreign gene expression can be detected by enzymatic or immunological assays (for example, immuno-precipitation, radioimmunoassay, or immunoblotting). Naturally occurring membrane glycoproteins produced from recombinant vaccinia infected cells
25 are glycosylated and may be transported to the cell surface. High expressing levels can be obtained by using strong promoters.

Other requirements for viral vectors for use in vaccines include good immunogenicity and safety. MVA is a replication-impaired vaccinia strain with a good safety record.
30 In most cell types and normal human tissue, MVA does not replicate. Replication of MVA is observed in a few transformed cell types such as BHK21 cells. Carroll *et al* (1997) have shown that the recombinant MVA is equally as good as conventional

recombinant vaccinia vectors at generating a protective CD8+T cell response and is an efficacious alternative to the more commonly used replication competent vaccinia virus. The vaccinia virus strains derived from MVA, or independently developed strains having the features of MVA which make MVA particularly suitable for use in a vaccine, are also suitable for use in the present invention.

Preferably, the vector is a vaccinia virus vector such as MVA or NYVAC. Most preferred is the vaccinia strain modified virus ankara (MVA) or a strain derived therefrom. Alternatives to vaccinia vectors include avipox vectors such as fowlpox or canarypox known as ALVAC and strains derived therefrom which can infect and express recombinant proteins in human cells but are unable to replicate.

In one aspect of the present invention at least one immune evasion gene is deleted from the poxvirus vector.

Viruses, especially large viruses such as poxviruses which have an extensive coding capacity and can thus encode a variety of genes, have developed a number of techniques for evading the immune system of their hosts. For example, they are able to evade non-specific defences such as complement, interferons and the inflammatory response, as well as to interfere with or block the function of cytokines. A number of these immune evasion polypeptides have been deleted from MVA, with the exception of the interferon resistance protein in the left terminal region.

Poxviruses in general, being large DNA viruses which establish acute, rather than latent, infections. They encode so many antigenic proteins that antigenic variation is difficult, thus relying on active immune evasion to protect themselves from the mammalian immune system. They possess a number of genes which encode polypeptides which are responsible for interfering with a number of aspects of the immune system: they disrupt interferon action, interfere with complement, cytokine activity, inflammatory responses and CTL recognition (for a review, Smith *et al.*, (1997) Immunol Rev 159:137-154). Removal of these proteins is beneficial in promoting the ability of weak immunogens encoded on a poxvirus vector to elicit an immune response in a subject.

5

10

15

20

25

30

The C21L gene is in turn related to C4b-binding protein in humans, and interacts with cells bearing C4b on the surface to prevent binding to the CR1 complement receptor.

Soluble Cytokine Receptors. The product of the vaccinia WR B15R gene (B16R in Copenhagen strain vaccinia) is related to IL1-R, the receptor for IL-1 β .

The WR gene ORF SalF19R, A53R in Copenhagen strain vaccinia, encodes a TNF receptor. However, in wild-type virus both of these genes are believed to be inactive due to fragmentation of the ORFs.

10

The B8R gene is believed to encode a soluble IFN- γ receptor, providing the virus with yet another IFN evasion mechanism.

Inflammation. A number of genes are believed to be involved in the prevention of inflammatory responses to viral infection. These include A44L, K2L, B13R and B22R.

In one aspect of the present invention, the majority of the immune evasion genes are deleted from the recombinant poxvirus vector. Preferably, all the immune evasion genes are deleted. Thus, in one aspect of the present invention, the recombinant poxvirus vector is a recombinant MVA vector in which the K3L interferon resistance protein gene has been disrupted or deleted.

Preferred are poxviruses which are non-hazardous to the intended subject. Thus, for example, for use in humans, poxviruses which are either host-range restricted, such as avipox viruses, or otherwise attenuated, such as attenuated strains of vaccinia (including NYVAC and MVA) are preferred. Most preferred are attenuated vaccinia virus strains, although non-vaccinia strains are usefully employed in subjects with pre-existing smallpox immunity.

30

A construct which contains at least one nucleic acid which codes for 5T4 flanked by MVA DNA sequences adjacent to a naturally occurring deletion, e.g. deletion II,

within the MVA genome, is introduced into cells infected with MVA, to allow homologous recombination.

Once the construct has been introduced into the eukaryotic cell and the 5T4 DNA has
 5 recombined with the viral DNA, the desired recombinant vaccinia virus, can be isolated, preferably with the aid of a marker (Nakano *et al* Proc. Natl. Acad. Sci. USA 79, 1593-1596 [1982], Franke *et al* Mol. Cell. Biol. 1918-1924 [1985], Chakrabarti *et al* Mol. Cell. Biol. 3403-3409 [1985], Fathi *et al* Virology 97-105 [1986]).

10 The construct to be inserted can be linear or circular. A circular DNA is preferred, especially a plasmid. The construct contains sequences flanking the left and the right side of a naturally occurring deletion, e.g. deletion II, within the MVA genome (Altenburger, W., Suter, C.P. and Altenburger J. (1989) Arch. Virol. 105, 15-27). The foreign DNA sequence is inserted between the sequences flanking the naturally
 15 occurring deletion.

For the expression of at least one nucleic acid, it is necessary for regulatory sequences, which are required for the transcription of the nucleic acid to be present upstream of the nucleic acid. Such regulatory sequences are known to those skilled in the art, and
 20 includes for example those of the vaccinia 11 kDa gene as are described in EP-A-198,328, and those of the 7.5 kDa gene (EP-A-110,385).

The construct can be introduced into the MVA infected cells by transfection, for example by means of calcium phosphate precipitation (Graham *et al* Virol. 52, 456-
 25 467 [1973; Wigler *et al* Cell 777-785 [1979] by means of electroporation (Neumann *et al* EMBO J. 1, 841-845 [1982]), by microinjection (Graessmann *et al* Meth. Enzymology 101, 482-492 (1983)), by means of liposomes (Straubinger *et al* Methods in Enzymology 101, 512-527 (1983)), by means of spheroplasts (Schaffner, Proc. Natl. Acad. Sci. USA 77, 2163-2167 (1980)) or by other methods known to those skilled in
 30 the art. Transfection by means of liposomes is preferred.

The recombinant priming and boosting vectors of the present invention can have a tropism for a specific cell type in the mammal. By way of example, the recombinant vectors of the present invention can be engineered to infect professional APCs such as dendritic cells and macrophages. Dendritic cells are known to be orchestrators of a successful immune response especially that of a cell mediated response. It has been shown that *ex vivo* treatment of dendritic cells with antigen or viral vectors containing such a target antigen, will induce efficacious immune responses when infused into syngeneic animals or humans (see Nestle FO, *et al.* Vaccination of melanoma patients with peptide- or tumor lysate-pulsed dendritic cells, Nat Med. 1998 Mar;4(3):328-32 and Kim CJ, *et al.* Dendritic cells infected with poxviruses encoding MART-1/Melan A sensitize T lymphocytes *in vitro*. J Immunother. 1997 Jul;20(4):276-86. The recombinant vectors can also infect tumour cells. Alternatively, the recombinant vectors are able to infect any cell in the mammal.

Other examples of vectors include *ex vivo* delivery systems, which include but are not limited to DNA transfection methods such as electroporation, DNA biolistics, lipid-mediated transfection and compacted DNA-mediated transfection.

The vector may be a plasmid DNA vector. As used herein, "plasmid" refers to discrete elements that are used to introduce heterologous DNA into cells for either expression or replication thereof. Selection and use of such vehicles are well within the skill of the artisan. Many plasmids are available, and selection of appropriate plasmid will depend on the intended use of the plasmid, i.e. whether it is to be used for DNA amplification or for DNA expression, the size of the DNA to be inserted into the plasmid, and the host cell to be transformed with the plasmid. Each plasmid contains various components depending on its function (amplification of DNA or expression of DNA) and the host cell for which it is compatible. The plasmid components generally include, but are not limited to, one or more of the following: an origin of replication, one or more marker genes, an enhancer element, a promoter, a transcription termination sequence and a signal sequence.

Both expression and cloning plasmids generally contain nucleic acid sequence that enable the plasmid to replicate in one or more selected host cells. Typically in cloning plasmids, this sequence is one that enables the plasmid to replicate independently of the host chromosomal DNA, and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria, yeast and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2m plasmid origin is suitable for yeast, and various viral origins (e.g. SV 40, polyoma, adenovirus) are useful for cloning plasmids in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression plasmids unless these are used in mammalian cells competent for high level DNA replication, such as COS cells.

Most expression plasmids are shuttle plasmids, i.e. they are capable of replication in at least one class of organisms but can be transfected into another class of organisms for expression. For example, a plasmid is cloned in *E. Coli* and then the same plasmid is transfected into yeast or mammalian cells even though it is not capable of replicating independently of the host cell chromosome.

Advantageously, an expression and cloning plasmid may contain a selection gene also referred to as selectable marker. This gene encodes a protein necessary for the survival or growth of transformed host cells grown in a selective culture medium. Host cells not transformed with the plasmid containing the selection gene will not survive in the culture medium. Typical selection genes encode proteins that confer resistance to antibiotics and other toxins, e.g. ampicillin, neomycin, methotrexate or tetracycline, complement auxotrophic deficiencies, or supply critical nutrients not available from complex media.

As to a selective gene marker appropriate for yeast, any marker gene can be used which facilitates the selection for transformants due to the phenotypic expression of the marker gene. Suitable markers for yeast are, for example, those conferring resistance to antibiotics G418, hygromycin or bleomycin, or provide for prototrophy in

an auxotrophic yeast mutant, for example the URA3, LEU2, LYS2, TRP1, or HIS3 gene.

- Since the replication of plasmids is conveniently done in *E. Coli*, an *E. Coli* genetic marker and an *E. Coli* origin of replication are advantageously included. These can be obtained from *E. Coli* plasmids, such as pBR322, Bluescript© plasmid or a pUC plasmid, e.g. pUC18 or pUC19, which contain both *E. Coli* replication origin and *E. Coli* genetic marker conferring resistance to antibiotics, such as ampicillin.
- 10 Suitable selectable markers for mammalian cells are those that enable the identification of cells which have taken up 5T4 nucleic acid, such as dihydrofolate reductase (DHFR, methotrexate resistance), thymidine kinase, or genes conferring resistance to G418 or hygromycin. The mammalian cell transformants are placed under selection pressure which only those transformants which have taken up and are expressing the marker are
- 15 uniquely adapted to survive. In the case of a DHFR or glutamine synthase (GS) marker, selection pressure can be imposed by culturing the transformants under conditions in which the pressure is progressively increased, thereby leading to amplification (at its chromosomal integration site) of both the selection gene and the linked DNA that encodes 5T4. Amplification is the process by which genes in greater
- 20 demand for the production of a protein critical for growth, together with closely associated genes which may encode a desired protein, are reiterated in tandem within the chromosomes of recombinant cells. Increased quantities of desired protein are usually synthesised from thus amplified DNA.
- 25 Expression and cloning plasmids usually contain a promoter that is recognised by the host organism and is operably linked to 5T4 nucleic acid. Such a promoter may be inducible or constitutive. The promoters are operably linked to DNA encoding 5T4 by removing the promoter from the source DNA by restriction enzyme digestion and inserting the isolated promoter sequence into the plasmid. Both the native 5T4
- 30 promoter sequence and many heterologous promoters may be used to direct amplification and/or expression of 5T4 DNA. The term "operably linked" refers to a juxtaposition wherein the components described are in a relationship permitting them

to function in their intended manner. A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

- 5 Promoters suitable for use with prokaryotic hosts include, for example, the β -lactamase and lactose promoter systems, alkaline phosphatase, the tryptophan (trp) promoter system and hybrid promoters such as the tac promoter. Their nucleotide sequences have been published, thereby enabling the skilled worker operably to ligate them to DNA encoding 5T4, using linkers or adaptors to supply any required
10 restriction sites. Promoters for use in bacterial systems will also generally contain a Shine-Delgarno sequence operably linked to the DNA encoding 5T4.

Preferred expression plasmids are bacterial expression plasmids which comprise a promoter of a bacteriophage such as phagex or T7 which is capable of functioning in
15 the bacteria. In one of the most widely used expression systems, the nucleic acid encoding the fusion protein may be transcribed from the plasmid by T7 RNA polymerase (Studier *et al*, Methods in Enzymol. 185; 60-89, 1990). In the *E. Coli* BL21(DE3) host strain, used in conjunction with pET plasmids, the T7 RNA polymerase is produced from the λ -lysogen DE3 in the host bacterium, and its
20 expression is under the control of the IPTG inducible lac UV5 promoter. This system has been employed successfully for over-production of many proteins. Alternatively the polymerase gene may be introduced on a lambda phage by infection with an int-phage such as the CE6 phage which is commercially available (Novagen, Madison, USA). other plasmids include plasmids containing the lambda PL promoter such as
25 PLEX (Invitrogen, NL) , plasmids containing the trc promoters such as pTrcHisXpressTm (Invitrogen) or pTrc99 (Pharmacia Biotech, SE) , or plasmids containing the tac promoter such as pKK223-3 (Pharmacia Biotech) or PMAL (new England Biolabs, MA, USA).

- 30 Moreover, the 5T4 gene according to the invention preferably includes a secretion sequence in order to facilitate secretion of the polypeptide from bacterial hosts, such that it will be produced as a soluble native peptide rather than in an inclusion body.

The peptide may be recovered from the bacterial periplasmic space, or the culture medium, as appropriate.

- Suitable promoting sequences for use with yeast hosts may be regulated or constitutive and are preferably derived from a highly expressed yeast gene, especially a *Saccharomyces cerevisiae* gene. Thus, the promoter of the TRP1 gene, the ADHI or ADHII gene, the acid phosphatase (PH05) gene, a promoter of the yeast mating pheromone genes coding for the a- or α -factor or a promoter derived from a gene encoding a glycolytic enzyme such as the promoter of the enolase, glyceraldehyde-3-phosphate dehydrogenase (GAP), 3-phospho glycerate kinase (PGK), hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triose phosphate isomerase, phosphoglucose isomerase or glucokinase genes, the *S. cerevisiae* GAL 4 gene, the *S. pombe* nmt 1 gene or a promoter from the TATA binding protein (TBP) gene can be used. Furthermore, it is possible to use hybrid promoters comprising upstream activation sequences (UAS) of one yeast gene and downstream promoter elements including a functional TATA box of another yeast gene, for example a hybrid promoter including the UAS(s) of the yeast PH05 gene and downstream promoter elements including a functional TATA box of the yeast GAP gene (PH05-GAP hybrid promoter). A suitable constitutive PH05 promoter is e.g. a shortened acid phosphatase PH05 promoter devoid of the upstream regulatory elements (UAS) such as the PH05 (-173) promoter element starting at nucleotide -173 and ending at nucleotide -9 of the PH05 gene.
- 5T4 gene transcription from plasmids in mammalian hosts may be controlled by promoters derived from the genomes of viruses such as polyoma virus, adenovirus, fowlpox virus, bovine papilloma virus, avian sarcoma virus, cytomegalovirus (CMV), a retrovirus and Simian Virus 40 (SV40), from heterologous mammalian promoters such as the actin promoter or a very strong promoter, e.g. a ribosomal protein promoter, and from the promoter normally associated with 5T4 sequence, provided such promoters are compatible with the host cell systems.

Transcription of a DNA encoding 5T4 by higher eukaryotes may be increased by inserting an enhancer sequence into the plasmid. Enhancers are relatively orientation and position independent. Many enhancer sequences are known from mammalian genes (e.g. elastase and globin). However, typically one will employ an enhancer from
 5 a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270) and the CMV early promoter enhancer. The enhancer may be spliced into the plasmid at a position 5' or 3' to 5T4 DNA, but is preferably located at a site 5' from the promoter.

10 Advantageously, a eukaryotic expression plasmid encoding 5T4 may comprise a locus control region (LCR). LCRs are capable of directing high-level integration site independent expression of transgenes integrated into host cell chromatin, which is of importance especially where the 5T4 gene is to be expressed in the context of a permanently-transfected eukaryotic cell line in which chromosomal integration of the
 15 plasmid has occurred, in plasmids designed for gene therapy applications or in transgenic animals.

Eukaryotic expression plasmids will also contain sequences necessary for the termination of transcription and for stabilising the mRNA. Such sequences are
 20 commonly available from the 5' and 3' untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding 5T4.

An expression plasmid includes any plasmid capable of expressing 5T4 nucleic acids
 25 that are operatively linked with regulatory sequences, such as promoter regions, that are capable of expression of such DNAs. Thus, an expression plasmid refers to a recombinant DNA or RNA construct, such as a plasmid, a phage, recombinant virus or other plasmid, that upon introduction into an appropriate host cell, results in expression of the cloned DNA. Appropriate expression plasmids are well known to
 30 those with ordinary skill in the art and include those that are replicable in eukaryotic and/or prokaryotic cells and those that remain episomal or those which integrate into the host cell genome. For example, DNAs encoding 5T4 may be inserted into a

plasmid suitable for expression of cDNAs in mammalian cells, e.g. a CMV enhancer-based plasmid such as pEVRF (Matthias, *et al.*, (1989) NAR 17, 6418).

Particularly useful for practising the present invention are expression plasmids that provide for the transient expression of DNA encoding 5T4 in mammalian cells. Transient expression usually involves the use of an expression plasmid that is able to replicate efficiently in a host cell, such that the host cell accumulates many copies of the expression plasmid, and, in turn, synthesises high levels of 5T4. For the purposes of the present invention, transient expression systems are useful e.g. for identifying 5T4 mutants, to identify potential phosphorylation sites, or to characterise functional domains of the protein.

Construction of plasmids according to the invention employs conventional ligation techniques. Isolated plasmids or DNA fragments are cleaved, tailored, and religated in the form desired to generate the plasmids required. If desired, analysis to confirm correct sequences in the constructed plasmids is performed in a known fashion. Suitable methods for constructing expression plasmids, preparing *in vitro* transcripts, introducing DNA into host cells, and performing analyses for assessing 5T4 expression and function are known to those skilled in the art. Gene presence, amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA, dot blotting (DNA or RNA analysis), or *in situ* hybridisation, using an appropriately labelled probe which may be based on a sequence provided herein. Those skilled in the art will readily envisage how these methods may be modified, if desired.

25

5T4 antigen, fragments and variants

5T4 antigen, as referred to herein, includes peptides and other fragments of 5T4 which retain at least one common antigenic determinant of 5T4.

30

"Common antigenic determinant" means that the derivative in question at least one antigenic function of 5T4. Antigenic functions includes possession of an epitope or

antigenic site that is capable of cross-reacting with antibodies raised against a naturally occurring or denatured 5T4 polypeptide or fragment thereof, or the ability to bind HLA molecules and induce a 5T4-specific immune response. Thus 5T4 as provided by the present invention includes splice variants encoded by mRNA generated by alternative splicing of a primary transcript, amino acid mutants, glycosylation variants and other covalent derivatives of 5T4 which retain the physiological and/or physical properties of 5T4. Exemplary derivatives include molecules wherein the protein of the invention is covalently modified by substitution, chemical, enzymatic, or other appropriate means with a moiety other than a naturally occurring amino acid. Such a moiety may be a detectable moiety such as an enzyme or a radioisotope. Further included are naturally occurring variants of 5T4 found with a particular species, preferably a mammal. Such a variant may be encoded by a related gene of the same gene family, by an allelic variant of a particular gene, or represent an alternative splicing variant of the 5T4 gene.

Derivatives which retain common antigenic determinants can be fragments of 5T4. Fragments of 5T4 comprise individual domains thereof, as well as smaller polypeptides derived from the domains. Preferably, smaller polypeptides derived from 5T4 according to the invention define a single epitope which is characteristic of 5T4. Fragments may in theory be almost any size, as long as they retain one characteristic of 5T4. Preferably, fragments will be between 5 and 400 amino acids in length. Longer fragments are regarded as truncations of the full-length 5T4 and generally encompassed by the term "5T4". Advantageously, fragments are relatively small peptides of the order of 5 to 25 amino acids in length. Preferred are peptides about 9 amino acids in length.

Derivatives of 5T4 also comprise mutants thereof, which may contain amino acid deletions, additions or substitutions, subject to the requirement to maintain at least one feature characteristic of 5T4. Thus, conservative amino acid substitutions may be made substantially without altering the nature of 5T4, as may truncations from the 5' or 3' ends. Deletions and substitutions may moreover be made to the fragments of 5T4 comprised by the invention. 5T4 mutants may be produced from a DNA encoding 5T4

which has been subjected to *in vitro* mutagenesis resulting e.g. in an addition, exchange and/or deletion of one or more amino acids. For example, substitutional, deletional or insertional variants of 5T4 can be prepared by recombinant methods and screened for immuno-crossreactivity with the native forms of 5T4.

5

Moreover, variant peptides can be screened for superior HLA binding capabilities using the program "Peptide Binding Predictions" devised by K. Parker at the National Institutes of Health (see Parker, K. C *et al.* 1994.J.Immunol. 152:163).

- 10 The fragments, mutants and other derivative of 5T4 preferably retain substantial homology with 5T4. As used herein, "homology" means that the two entities share sufficient characteristics for the skilled person to determine that they are similar in origin and function. Preferably, homology is used to refer to sequence identity. Thus, the derivatives of 5T4 preferably retain substantial sequence identity with the sequence
- 15 of SEQ ID No. 2.

- "Substantial homology", where homology indicates sequence identity, means more than 40% sequence identity, preferably more than 45% sequence identity and most preferably a sequence identity of 50% or more, as judged by direct sequence alignment
- 20 and comparison.

- Sequence homology (or identity) may moreover be determined using any suitable homology algorithm, using for example default parameters. Advantageously, the BLAST algorithm is employed, with parameters set to default values. The BLAST
- 25 algorithm is described in detail at http://www.ncbi.nih.gov/BLAST/blast_help.html, which is incorporated herein by reference. The search parameters are defined as follows, and are advantageously set to the defined default parameters.

- Advantageously, "substantial homology" when assessed by BLAST equates to
- 30 sequences which match with an EXPECT value of at least about 7, preferably at least about 9 and most preferably 10 or more. The default threshold for EXPECT in BLAST searching is usually 10.

BLAST (Basic Local Alignment Search Tool) is the heuristic search algorithm employed by the programs `blastp`, `blastn`, `blastx`, `tblastn`, and `tblastx`; these programs ascribe significance to their findings using the statistical methods of Karlin and Altschul (see http://www.ncbi.nih.gov/BLAST/blast_help.html) with a few enhancements. The BLAST programs were tailored for sequence similarity searching, for example to identify homologues to a query sequence. The programs are not generally useful for motif-style searching. For a discussion of basic issues in similarity searching of sequence databases, see Altschul *et al.* (1994) *Nature Genetics* 6:119-129.

10

The five BLAST programs available at <http://www.ncbi.nlm.nih.gov> perform the following tasks:

15

blastp compares an amino acid query sequence against a protein sequence database;

blastn compares a nucleotide query sequence against a nucleotide sequence database;

20

blastx compares the six-frame conceptual translation products of a nucleotide query sequence (both strands) against a protein sequence database;

tblastn compares a protein query sequence against a nucleotide sequence database dynamically translated in all six reading frames (both strands).

25

tblastx compares the six-frame translations of a nucleotide query sequence against the six-frame translations of a nucleotide sequence database.

BLAST uses the following search parameters:

30

HISTOGRAM Display a histogram of scores for each search; default is yes. (See parameter H in the BLAST Manual).

DESCRIPTIONS Restricts the number of short descriptions of matching sequences reported to the number specified; default limit is 100 descriptions. (See parameter V in the manual page). See also EXPECT and CUTOFF.

- 5 ALIGNMENTS Restricts database sequences to the number specified for which high-scoring segment pairs (HSPs) are reported; the default limit is 50. If more database sequences than this happen to satisfy the statistical significance threshold for reporting (see EXPECT and CUTOFF below), only the matches ascribed the greatest statistical significance are reported. (See parameter B in the BLAST Manual).

10

EXPECT The statistical significance threshold for reporting matches against database sequences; the default value is 10, such that 10 matches are expected to be found merely by chance, according to the stochastic model of Karlin and Altschul (1990). If the statistical significance ascribed to a match is greater than the EXPECT threshold,
15 the match will not be reported. Lower EXPECT thresholds are more stringent, leading to fewer chance matches being reported. Fractional values are acceptable. (See parameter E in the BLAST Manual).

20

CUTOFF Cutoff score for reporting high-scoring segment pairs. The default value is calculated from the EXPECT value (see above). HSPs are reported for a database sequence only if the statistical significance ascribed to them is at least as high as would be ascribed to a lone HSP having a score equal to the CUTOFF value. Higher CUTOFF values are more stringent, leading to fewer chance matches being reported. (See parameter S in the BLAST Manual). Typically, significance thresholds can be
25 more intuitively managed using EXPECT.

- MATRIX Specify an alternate scoring matrix for BLASTP, BLASTX, TBLASTN and TBLASTX. The default matrix is BLOSUM62 (Henikoff & Henikoff, 1992). The valid alternative choices include: PAM40, PAM120, PAM250 and IDENTITY. No
30 alternate scoring matrices are available for BLASTN; specifying the MATRIX directive in BLASTN requests returns an error response.

5 FILTER Mask off segments of the query sequence that have low compositional complexity, as determined by the SEG program of Wootton & Federhen (1993) Computers and Chemistry 17:149-163, or segments consisting of short-periodicity internal repeats, as determined by the XNU program of Claverie & States (1993) Computers and Chemistry 17:191-201, or, for BLASTN, by the DUST program of Tatusov and Lipman (see <http://www.ncbi.nlm.nih.gov>). Filtering can eliminate statistically significant but biologically uninteresting reports from the blast output (e.g., hits against common acidic-, basic- or proline-rich regions), leaving the more biologically interesting regions of the query sequence available for specific matching against database sequences.

Low complexity sequence found by a filter program is substituted using the letter "N" in nucleotide sequence (e.g., "NNNNNNNNNNNNNNNN") and the letter "X" in protein sequences (e.g., "XXXXXXXXXX").

It is not unusual for nothing at all to be masked by SEG, XNU, or both, when applied to sequences in SWISS-PROT, so filtering should not be expected to always yield an effect. Furthermore, in some cases, sequences are masked in their entirety, indicating that the statistical significance of any matches reported against the unfiltered query sequence should be suspect.

NCBI-gi Causes NCBI gi identifiers to be shown in the output, in addition to the
30 accession and/or locus name.

Most preferably, sequence comparisons are conducted using the simple BLAST search algorithm provided at <http://www.ncbi.nlm.nih.gov/BLAST>.

Alternatively, sequence homology may be determined by algorithms such as FastA,
5 available at <http://biology.ncsa.uiuc.edu/BW30/BW.cgi>. FastA is considered to be superior to BLAST for alignment of short sequences. Advantageously, the FastA algorithm is employed using default parameters at <http://biology.ncsa.uiuc.edu/BW30/BW.cgi>.

10 Preferably, the protein or derivative thereof of the invention is provided in isolated form. "Isolated" means that the protein or derivative has been identified and is free of one or more components of its natural environment. Isolated 5T4 includes 5T4 in a recombinant cell culture. 5T4 present in an organism expressing a recombinant 5T4 gene, whether the 5T4 protein is "isolated" or otherwise, is included within the scope
15 of the present invention.

In the vaccination of humans against tumours, the use of non-human TAAs is preferred. The invention accordingly provides canine 5T4. Canine 5T4 is advantageously provided for use as a vaccine component in humans, in order to elicit
20 an immune response to human 5T4 in a human subject.

The sequence of canine 5T4 is set forth in SEQ. ID. No. 3. Sequences from other canine sources are obtainable by those skilled in the art, for example by hybridisation with a nucleic acid probe derived from SEQ. ID. No. 3.

25 Exemplary nucleic acids can thus be characterised as those nucleotide sequences which encode a canine 5T4 protein and hybridise to the DNA sequences set forth SEQ ID No. 3, or a selected fragment of said DNA sequence. Preferred are such sequences encoding canine 5T4 which hybridise under high-stringency conditions to the
30 sequence of SEQ ID No. 3.

Stringency of hybridisation refers to conditions under which polynucleic acid hybrids are stable. Such conditions are evident to those of ordinary skill in the field. As known to those of skill in the art, the stability of hybrids is reflected in the melting temperature (T_m) of the hybrid which decreases approximately 1 to 1.5°C with every 1% decrease in sequence homology. In general, the stability of a hybrid is a function of sodium ion concentration and temperature. Typically, the hybridisation reaction is performed under conditions of higher stringency, followed by washes of varying stringency.

As used herein, high stringency refers to conditions that permit hybridisation of only those nucleic acid sequences that form stable hybrids in 1 M Na⁺ at 65-68 °C. High stringency conditions can be provided, for example, by hybridisation in an aqueous solution containing 6x SSC, 5x Denhardt's, 1 % SDS (sodium dodecyl sulphate), 0.1 Na⁺ pyrophosphate and 0.1 mg/ml denatured salmon sperm DNA as non specific competitor. Following hybridisation, high stringency washing may be done in several steps, with a final wash (about 30 min) at the hybridisation temperature in 0.2 - 0.1x SSC, 0.1 % SDS.

Moderate stringency refers to conditions equivalent to hybridisation in the above described solution but at about 60-62°C. In that case the final wash is performed at the hybridisation temperature in 1x SSC, 0.1 % SDS.

Low stringency refers to conditions equivalent to hybridisation in the above described solution at about 50-52°C. In that case, the final wash is performed at the hybridisation temperature in 2x SSC, 0.1 % SDS.

It is understood that these conditions may be adapted and duplicated using a variety of buffers, e.g. formamide-based buffers, and temperatures. Denhardt's solution and SSC are well known to those of skill in the art as are other suitable hybridisation buffers (see, e.g. Sambrook, *et al.*, eds. (1989) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, New York or Ausubel, *et al.*, eds. (1990) *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc.). Optimal

hybridisation conditions have to be determined empirically, as the length and the GC content of the probe also play a role.

Advantageously, the invention moreover provides nucleic acid sequence which are
5 capable of hybridising, under stringent conditions, to a fragment of SEQ. ID. No. 3. Preferably, the fragment is between 15 and 50 bases in length. Advantageously, it is about 25 bases in length.

Given the guidance provided herein, the nucleic acids of the invention are obtainable
10 according to methods well known in the art. For example, a DNA of the invention is obtainable by chemical synthesis, using polymerase chain reaction (PCR) or by screening a genomic library or a suitable cDNA library prepared from a source believed to possess canine 5T4 and to express it at a detectable level.

15 Chemical methods for synthesis of a nucleic acid of interest are known in the art and include triester, phosphite, phosphoramidite and H-phosphonate methods, PCR and other autoprimer methods as well as oligonucleotide synthesis on solid supports. These methods may be used if the entire nucleic acid sequence of the nucleic acid is known, or the sequence of the nucleic acid complementary to the coding strand is available.
20 Alternatively, if the target amino acid sequence is known, one may infer potential nucleic acid sequences using known and preferred coding residues for each amino acid residue.

An alternative means to isolate the gene encoding canine 5T4 is to use PCR
25 technology as described e.g. in section 14 of Sambrook *et al.*, 1989. This method requires the use of oligonucleotide probes that will hybridise to canine 5T4 nucleic acid. Strategies for selection of oligonucleotides are described below.

Libraries are screened with probes or analytical tools designed to identify the gene of
30 interest or the protein encoded by it. For cDNA expression libraries suitable means include monoclonal or polyclonal antibodies that recognise and specifically bind to canine 5T4; oligonucleotides of about 20 to 80 bases in length that encode known or

suspected canine 5T4 cDNA from the same or different species; and/or complementary or homologous cDNAs or fragments thereof that encode the same or a hybridising gene. Appropriate probes for screening genomic DNA libraries include, but are not limited to oligonucleotides, cDNAs or fragments thereof that encode the same or hybridising DNA; and/or homologous genomic DNAs or fragments thereof.

A nucleic acid encoding canine 5T4 may be isolated by screening suitable cDNA or genomic libraries under suitable hybridisation conditions with a probe, i.e. a nucleic acid disclosed herein including oligonucleotides derivable from the sequences set forth in SEQ ID NO. 3. Suitable libraries are commercially available or can be prepared e.g. from cell lines, tissue samples, and the like.

As used herein, a probe is e.g. a single-stranded DNA or RNA that has a sequence of nucleotides that includes between 10 and 50, preferably between 15 and 30 and most preferably at least about 20 contiguous bases that are the same as (or the complement of) an equivalent or greater number of contiguous bases set forth in SEQ ID No. 3. The nucleic acid sequences selected as probes should be of sufficient length and sufficiently unambiguous so that false positive results are minimised. The nucleotide sequences are usually based on conserved or highly homologous nucleotide sequences or regions of canine 5T4. The nucleic acids used as probes may be degenerate at one or more positions. The use of degenerate oligonucleotides may be of particular importance where a library is screened from a species in which preferential codon usage in that species is not known.

Preferred regions from which to construct probes include 5' and/or 3' coding sequences, sequences predicted to encode ligand binding sites, and the like. For example, either the full-length cDNA clone disclosed herein or fragments thereof can be used as probes. Preferably, nucleic acid probes of the invention are labelled with suitable label means for ready detection upon hybridisation. For example, a suitable label means is a radiolabel. The preferred method of labelling a DNA fragment is by incorporating $\alpha^{32}\text{P}$ dATP with the Klenow fragment of DNA polymerase in a random priming reaction, as is well known in the art. Oligonucleotides are usually end-labelled with $\gamma^{32}\text{P}$ -labelled ATP and polynucleotide kinase. However, other methods (e.g. non-

radioactive) may also be used to label the fragment or oligonucleotide, including e.g. enzyme labelling, fluorescent labelling with suitable fluorophores and biotinylation.

After screening the library, e.g. with a portion of DNA including substantially the entire canine 5T4-encoding sequence or a suitable oligonucleotide based on a portion of said DNA, positive clones are identified by detecting a hybridisation signal; the identified clones are characterised by restriction enzyme mapping and/or DNA sequence analysis, and then examined, e.g. by comparison with the sequences set forth herein, to ascertain whether they include DNA encoding a complete canine 5T4 (i.e., if they include translation initiation and termination codons). If the selected clones are incomplete, they may be used to rescreen the same or a different library to obtain overlapping clones. If the library is genomic, then the overlapping clones may include exons and introns. If the library is a cDNA library, then the overlapping clones will include an open reading frame. In both instances, complete clones may be identified by comparison with the DNAs and deduced amino acid sequences provided herein.

It is envisaged that the nucleic acid of the invention can be readily modified by nucleotide substitution, nucleotide deletion, nucleotide insertion or inversion of a nucleotide stretch, and any combination thereof. Such mutants can be used e.g. to produce a canine 5T4 mutant that has an amino acid sequence differing from the canine 5T4 sequences as found in nature. Mutagenesis may be predetermined (site-specific) or random. A mutation which is not a silent mutation must not place sequences out of reading frames and preferably will not create complementary regions that could hybridise to produce secondary mRNA structure such as loops or hairpins.

25

The foregoing considerations may also be applied to the isolation of alternative murine (SEQ. ID. No. 2) or human (SEQ. ID. No. 1) 5T4 antigens.

Administration of Vectors encoding 5T4

30

A pharmaceutical composition according to the invention is a composition of matter comprising a vector encoding a 5T4 antigen, as described, as an active ingredient. The

active ingredients of a pharmaceutical composition comprising the active ingredient according to the invention are contemplated to exhibit excellent therapeutic and/or prophylactic activity, for example, in the treatment and/or prophylaxis of tumours or other diseases associated with cell proliferation, infections and inflammatory conditions, when administered in amount which depends on the particular case. Dosage regima may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation.

10 The active compound may be administered in a convenient manner such as by the oral, intravenous (where water soluble), intramuscular, subcutaneous, intranasal, intradermal or suppository routes or implanting (e.g. using slow release molecules). Depending on the route of administration, the active ingredient may be required to be coated in a material to protect said ingredients from the action of enzymes, acids and
15 other natural conditions which may inactivate said ingredient.

In order to administer the active compound by other than parenteral administration, it will be coated by, or administered with, a material to prevent its inactivation. For example, the active compound may be administered in an adjuvant, co-administered
20 with enzyme inhibitors or in liposomes. Adjuvant is used in its broadest sense and includes any immune stimulating compound such as interferon. Adjuvants contemplated herein include resorcinols, non-ionic surfactants such as polyoxyethylene oleyl ether and n-hexadecyl polyethylene ether. Enzyme inhibitors include pancreatic trypsin.

25 Liposomes include water-in-oil-in-water CGF emulsions as well as conventional liposomes.

The active compound may also be administered parenterally or intraperitoneally.
30 Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. In all cases the form must be
5 sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like),
10 suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants.

The prevention of the action of microorganisms can be brought about by various
15 antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminium monostearate and gelatin.

20 Sterile injectable solutions are prepared by incorporating the active compound in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilisation. Generally, dispersions are prepared by incorporating the sterilised active ingredient into a sterile
25 vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze-drying technique which yield a powder of the active ingredient plus any additional desired ingredient from previously sterile-filtered solution thereof.

30 When the active compound is suitably protected as described above, it may be orally administered, for example, with an inert diluent or with an assimilable edible carrier,

or it may be enclosed in hard or soft shell gelatin capsules, or it may be compressed into tablets, or it may be incorporated directly with the food of the diet. For oral therapeutic administration, the active compound may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. The amount of active compound in such therapeutically useful compositions is such that a suitable dosage will be obtained.

The tablets, troches, pills, capsules and the like may also contain the following: a binder such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin may be added or a flavouring agent such as peppermint, oil of wintergreen, or cherry flavouring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier.

Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the active compound, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavouring such as cherry or orange flavour. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compound may be incorporated into sustained-release preparations and formulations.

As used herein "pharmaceutically acceptable carrier and/or diluent" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, use thereof in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active material
 5 calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the novel dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active material and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such as active material for the treatment
 10 of disease in living subjects having a diseased condition in which bodily health is impaired.

The principal active ingredients are compounded for convenient and effective administration in effective amounts with a suitable pharmaceutically acceptable carrier
 15 in dosage unit form. In the case of compositions containing supplementary active ingredients, the dosages are determined by reference to the usual dose and manner of administration of the said ingredients.

Regimes for administration of 5T4-expressing vectors according to the present
 20 invention may be determined by conventional efficacy testing. Especially preferred, however, are regimes which include successive priming and boosting steps. It is observed that such regimes achieve superior breaking of immune tolerance and induction of CTL responses. In a preferred embodiment, the priming step is undertaken using a non-viral vector, such as a plasmid encoding 5T4, whilst boosting
 25 is undertaken using a viral vector, such as a poxvirus vector, encoding 5T4 (see Schneider *et al.*, 1998 Nat Med 4:397-402).

Administration of 5T4 antigen

In general, approaches outlined above relating to the administration of 5T4-encoding nucleic acids may be used for the administration of 5T4 antigen, as a conventional vaccine preparation, for the therapy and/or prophylaxis of tumours.

- 5 In general, vaccines may be prepared 5T4 antigen. The preparation of vaccines which contain an 5T4 as active ingredient(s) is known to one skilled in the art. Typically, such vaccines are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection may also be prepared. The preparation may also be emulsified, or the protein encapsulated in liposomes. The active immunogenic ingredients are often mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof.
- 10 In addition, if desired, the vaccine may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, and/or adjuvants which enhance the effectiveness of the vaccine. Examples of adjuvants which may be effective include but are not limited to: aluminium hydroxide, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-nor-muramyl-L-alanyl-D-isoglutamine (CGP 11637, referred to as nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (CGP 19835A, referred to as MTP-PE), and RIBI, which contains three components extracted from bacteria, monophosphoryl lipid A, trehalose dimycolate and cell wall skeleton (MPL+TDM+CWS) in a 2% squalene/Tween 80 emulsion.
- 25 Further examples of adjuvants and other agents include aluminium hydroxide, aluminium phosphate, aluminium potassium sulphate (alum), beryllium sulphate, silica, kaolin, carbon, water-in-oil emulsions, oil-in-water emulsions, muramyl dipeptide, bacterial endotoxin, lipid X, *Corynebacterium parvum* (*Propionobacterium acnes*), *Bordetella pertussis*, polyribonucleotides, sodium alginate, lanolin, lysolecithin, vitamin A, saponin, liposomes, levamisole, DEAE-dextran, blocked copolymers or other synthetic adjuvants. Such adjuvants are available commercially
- 30

from various sources, for example, Merck Adjuvant 65 (Merck and Company, Inc., Rahway, N.J.) or Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, Michigan).

Typically, adjuvants such as Amphigen (oil-in-water), Alhydrogel (aluminium hydroxide), or a mixture of Amphigen and Alhydrogel are used. Only aluminium hydroxide is approved for human use.

The proportion of immunogen and adjuvant can be varied over a broad range so long as both are present in effective amounts. For example, aluminium hydroxide can be present in an amount of about 0.5% of the vaccine mixture (Al_2O_3 basis). Conveniently, the vaccines are formulated to contain a final concentration of immunogen in the range of from 0.2 to 200 $\mu\text{g/ml}$, preferably 5 to 50 $\mu\text{g/ml}$, most preferably 15 $\mu\text{g/ml}$.

After formulation, the vaccine may be incorporated into a sterile container which is then sealed and stored at a low temperature, for example 4°C , or it may be freeze-dried. Lyophilisation permits long-term storage in a stabilised form.

The vaccines are conventionally administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Additional formulations which are suitable for other modes of administration include suppositories and, in some cases, oral formulations. For suppositories, traditional binders and carriers may include, for example, polyalkylene glycols or triglycerides; such suppositories may be formed from mixtures containing the active ingredient in the range of 0.5% to 10%, preferably 1% to 2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain 10% to 95% of active ingredient, preferably 25% to 70%. Where the vaccine composition is lyophilised, the lyophilised material may be reconstituted prior to administration, e.g. as a suspension. Reconstitution is preferably effected in buffer

Capsules, tablets and pills for oral administration to a patient may be provided with an enteric coating comprising, for example, Eudragit "S", Eudragit "L", cellulose acetate, cellulose acetate phthalate or hydroxypropylmethyl cellulose.

5

5T4 may be formulated into the vaccine as neutral or salt forms. Pharmaceutically acceptable salts include the acid addition salts (formed with free amino groups of the peptide) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids such as acetic, oxalic, tartaric and maleic.

10 Salts formed with the free carboxyl groups may also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine and procaine.

15 Measurement of the efficacy of 5T4 administration

5T4 activity as an immunotherapeutic molecule may be assessed according to any techniques known in the art, including assays for antibody production, induction of CTL responses and tumour regression models. Exemplary techniques are set forth in
20 the following examples.

The invention is further described, for the purposes of illustration only, in the following examples in which reference is made to the following Figures.

25 Figure 1a shows a gene construct;

Figure 1b shows a gene construct;

Figure 2a shows a photographic representation;

30

Figure 2b shows a photographic representation;

Figure 3a present a graph;

Figure 3b present a graph;

5 Figure 4a presents a graph;

Figure 4b presents a graph;

Figure 4c presents a graph;

10

Figure 5 presents a graph;

Figure 6 presents a graph;

Figure 7 presents a graph;

15

Figure 8 presents a graph; and

Figure 9 presents a graph;

20 In slightly more detail:

Figure 1a is a map of recombinant vaccinia virus MVA. Transgenes are placed under the control of the vaccinia virus synthetic early/late promoter. The *Lac Z* gene is under control of the vaccinia virus 7.5k early/late promoter. The DNA regions flanking these
25 genes are derived from the deleted region two of MVA, thus allowing homologous recombination into this site.

Figure 1b is a map of the recombinant vaccinia virus WR. The transgenes are under the control of the vaccinia virus synthetic early/late promoter. The *Lac Z* gene is under
30 control of the vaccinia virus 7.5k early/late promoter. The DNA regions flanking the genes are derived from the thymidine kinase (tk) gene of Wyeth strain VV, thus allowing recombination into this site.

Figure 2 shows a western blot of recombinant vaccinia virus expressing human 5T4. Samples are run on a 12% SDS PAGE and transferred to a nitrocellulose membrane.

- 5 Figure 2a: The blot is probed with a 1:500 dilution of MAb 5T4 (anti-human 5T4). Bound antibody is visualised with a anti-mouse HRP conjugated antibody and ECL. Lane 1: recombinant WR clone 1 expressing human 5T4; Lane 2: recombinant WR clone 2 expressing human 5T4; Lane 3: BS-C-1 cells infected with WR; Lane 4: uninfected BS-C-1 cells; Lane 5: B16 melanoma cells; Lane 6: B16 cell line
10 expressing human 5T4.

- Figure 2b: The blot is probed with a 1:500 dilution of rabbit anti-mouse 5T4. Lane 1: recombinant WR expressing LacZ; Lane 2: recombinant WR expressing mouse 5T4; lane 3: recombinant WR expressing human 5T4; lane 4: recombinant MVA expressing
15 mouse 5T4.

Figures 3a and 3b are graphs which show that inoculation of mice with MVA-h5T4 Protects against Challenge with CT26 expressing h5T4

- Figure 4a is a graph which shows that inoculation with MVA-h5T4 induces anti-
20 tumour activity against B16 tumours expressing h5T4.

Figures 4b and 4c are graphs which show that inoculation with MVA-m5T4 induces anti-tumour activity against B16 tumours expressing m5T4.

- 25 Figure 5 is a graph which shows that MVA-h5T4 induces tumour therapy in mice with pre-established lung tumours.

- Figure 6 is a graph which shows that mice that are vaccinated with MVA-m5T4 develop tumours at a slower rate than those that receive the control vaccine.
30

Figure 7 is a graph which shows that mice that were vaccinated with MVA-m5T4 had a lower tumour burden than those mice that received MVA-LacZ treatment.

EXAMPLES

Example 1

5

Construction of recombinant poxvirus vectors

Propagation of Vaccinia Virus

- 10 The highly attenuated strain MVA is derived from the replication competent strain Ankara and has endured over 570 passages in primary chick embryo fibroblast cells. MVA replication was initially thought to be restricted to CEF cells as only minimal replication in mammalian cells was reported. However, further analysis has shown that Baby Hamster Kidney cells (BHK-21) are able to support high titre production of
- 15 MVA. MVA may thus be grown on BHK-21 or primary CEF cells (Carroll & Moss (1997) Virology 238:198-211).

- To prepare CEF cells, 10 day old chick embryos are gutted and limbs and head are removed before being minced and trypsinised in a solution of 0.25% trypsin and
- 20 incubation at 37°C . The cell suspension is filtered through a course filter mesh and cells are washed and concentrated by centrifugation at 2000 rpm in a Sorvall RC-3B at 1500 rpm for 5 mins. Cells are suspended in MEM containing 10% FCS, aliquotted into 175cm flasks and incubated at 37°C in a CO₂ incubator. When monolayers are 95% confluent they are trypsinised and used to seed additional flasks or six well plates.
- 25 Alternatively, primary cultures are transferred to a 31°C incubator for later use (Sutter and Moss (1992) Proc Natl Acad Sci U S A 89:10847-10851).

Preparation of crude, semi-purified and purified virus stocks

- 30 Crude virus stocks are prepared for initial recombinant virus analysis or as viral stocks used for subsequent high titre virus preparations. Vaccinia virus preparations can be semi-purified by centrifuging out cell membranes and nuclei or by additional steps

involving sucrose centrifugation to prevent contamination by pre-expressed recombinant protein products and cellular organelles. Methods used are a modification of those described by Earl *et al.*, in: Ausubel *et al.* (Eds.), (1991) Current Protocols in Molecular Biology, pp. 16.16.1-16.16.17, New York: Greene Publishing Associates and Wiley Interscience; Earl and Moss, *ibid*, pp. 16.17.1-16.17.16; Earl and Moss, *ibid*, pp. 16.18.1-16.18.10; and Bronte *et al.*, (1997) Proc Natl Acad Sci U S A 94(7):3183-3188.

Crude Virus

10

MVA is grown in either CEF or BHK-21 (obtained from the ATCC) and WR is grown in HeLa or BS-C-1 (ATCC) in 175cm² tissue culture flasks. Briefly, confluent monolayers are infected with an moi of approx. 1 pfu with MVA or WR. Virus is suspended in 10ml MEM containing 2% FCS and added to 175cm² flasks containing
 15 confluent cell monolayers. After inoculation for 1 hour at 37°C an additional 20ml MEM containing 2% FCS is added. After 48-72 hours infected cells are scraped into the medium and pelleted at 1500g for 5 mins. For crude virus preparations cells are resuspended 2ml MEM (2%) per 175cm² flask. Cells are freeze thawed three times, sonicated and aliquotted into 1 ml freezing tubes. A representative aliquot is freeze
 20 thawed and titred to determine virus concentration. Virus stocks are stored below -20°C.

Semi-pure preparations

25 Infected cells are harvested as described previously (Earl *et al.*; Earl and Moss; 1991). After centrifugation cells are resuspended in PBS (2ml/175cm² flask) and homogenised by 30-40 strokes in a tight fitting glass dounce homogeniser, on ice. Cell breakage is checked by microscopy. Nuclei, cellular organelles and membranes are removed by a centrifugation at 300g for 5 mins (4°C), keep supernatant. The cell pellet
 30 is resuspended in 1ml/175cm² flask and centrifugation repeated. The supernatants are pooled, aliquoted and stored.

Purified preparation

Infected cells are harvested as previously described (Earl *et al.*; Earl and Moss; 1991) and resuspended in 10 mM Tris.Cl, pH 9.0 (2ml/flask), keeping samples on ice from this point of the procedure. Homogenise as described previously using 10 mM Tris. The lysate is sonicated (on ice) using an XL 2015 sonicating cup (Misonics, USA) at maximum output (500 W) for 1 min. The sample is placed on ice for 1 min and the sonication repeated up to 3 times. A maximum of 5ml is sonicated at a time, and ice is replenished during sonication. The lysate is gently layered onto a cushion of 17 ml of 36% sucrose (in 10 mM Tris.Cl, pH 9.0) in a SW-27 centrifuge tube. Lyates are centrifuged for 80 mins in an SW-27 rotor at 13 500 rpm (32,900 x g), 4°C. The supernatant is discarded and the viral pellet resuspended in sterile PBS and sonicated in a cup sonicator for 1 min (on ice). Concentrated virus is aliquoted and stored at below -20°C.

15

Example 2

Construction and Characterisation of Recombinant Virus vectors Expressing 5T4

Murine and human 5T4 genes are cloned into WR (pSC65) (Chakrabarti *et al.*, (1997) Biotechniques 23:1094-7) and MVA (pLW22) transfer plasmids to allow homologous recombination into targeted regions of the respective viral genomes.

Recombinant MVA and WR expressing human and murine 5T4

The 1.4kb murine and human 5T4 (supplied by P. Stern Paterson Institute Manchester) genes are excised from pBSII-m5T4 (pBluescript (Stratagene) containing the 5T4 cDNA) and pBSII-h5T4 (Myers *et al.*, (1994) JBC 269:9319-9324) respectively by *Eco* RI and *Bam* HI restriction digestion. The fragments are blunt ended by "filling in" with dNTPs and DNA polymerase. The blunt ended fragments are cloned into the *Pme*I site of pLW22 (an MVA transfer plasmid, consisting of an early late promoter (Chakrabarti *et al.*, 1997) upstream of an MCS. Adjacent is a VV 7.5Kb LacZ cassette, for detection of recombinant virus; see figure 1a), and *Sma* I site of pSC65 (b)

(Chakrabarti *et al.*, 1997). pLW22 and pSC65 direct homologous recombination to deletion region II and the tk gene respectively.

In constructs destined for vaccination of human subjects, the LacZ gene under the control of the 7.5k promoter is omitted. Recombinant plaques are identified by live immunostaining using an anti-5T4 monoclonal antibody, as described previously (Wyatt *et al.*, (1996) Vaccine 14:1451-1458).

Wild type MVA, supplied by B. Moss (NIH, Bethesda, USA) is grown in CEF cells from a plaque purified clone and is the same isolate that was used to make previously described recombinant viruses (Sutter and Moss (1992) Proc Natl Acad Sci U S A 89:10847-10851, Sutter *et al* (1994) Vaccine 12:1032-1040, Hirsch *et al* (1996) J Virol 70:3741-3752, Carroll and Moss (1995) Biotechniques 19: 352-355, Wyatt *et al* (1995) Virology 210:202-205, Sutter *et al.*, (1995) FEBS Lett. 371:9-12, Wyatt *et al.* (1996) Vaccine 14, 1451-1458, Carroll *et al* (1997) Vaccine, 15:387-394, Carroll and Moss (1997) Virology 238:198-211) The WR stock is supplied by B. Moss (NIH), from the ATCC isolate (see e.g. Earl and Moss, 1991). The WR stock is prepared in HeLa S3 cells (ATCC).

The protocol used to make recombinant MVA virus is similar to that described previously (Carroll & Moss (1997) Virology 238:198-211). Briefly: BHK-21 or CEF cells are infected at an moi of 0.1 with an MVA stock. Plasmid DNA is diluted to 2µg in 100ul d.H₂O and mixed with 30µg lipofectin (BRL) diluted to 100µl with sterile d.H₂O. After 10 minutes incubation at RT the Lipofectin/DNA solution is added to infected cells overlaid with Opti MEM. Five hours after incubation at 37°C cells media is aspirated and replaced with MEM containing 2% FCS. Cells are harvested after a further 36 hours incubation and assayed for the expression of β-gal on CEF or BHK-21 cells in the presence of 5-bromo-3-indolyl-D-galactosidase. Isolated plaques are plaque purified at least an additional 3 times. After plaque purification small viral stocks are prepared in CEF or BHK-21 cells.

Protocols for the construction of recombinant WR are similar to those described previously (Carroll and Moss (1995) Biotechniques 19: 352-355; Earl and Moss,

1991). Briefly: recombination is carried out as for MVA. However, BS-C-1 cells are used and recombinant plaques assayed in 143B tk⁻ cells in the presence of BrdU with an agar overlay containing the substrate for the LacZ gene, 5-bromo-3-indolyl-D-galactosidase. Neutral red is used to detect *LAC Z* negative spontaneous tk⁻ virus for
 5 evaluation of virus homogeneity.

Recombinant protein expression is initially analysed by direct plaque immunostaining using antibodies specific for h5T4 and m5T4 using a method similar to that described previously (Carroll & Moss (1997) *Virology* 238:198-211). Briefly: recombinant
 10 viruses are plaqued on monolayers of BS-C-1 cells, fixed with acetone/methanol and treated with MAb 5T4 (Hole N, and Stern PL, (1990) *Int J Cancer* 45(1):179-184). Anti-mouse HRP conjugated antibody and dianizidine substrate are used to visualise recombinant 5T4 protein expression. 5T4 expressed protein is further characterised by western blotting under non-reducing conditions, as the MAb recognises a
 15 conformational epitope. As can be seen in Figure 2 recombinant viruses express high levels of protein at the appropriate size of 72 kDa. The stock is checked for homogeneity by double-immunostaining as described in Carroll & Moss (1997) *Virology* 238:198-211.

20 **Example 3**

Animal Models to illustrate immunological cross protection of mouse 5T4 with human 5T4.

25 To determine if the 5T4 gene product from one species can induce immunity to 5T4 in another species, the recombinant poxviruses are tested in murine tumour models. The mouse models are based on CT26, a chemically induced adenocarcinoma of BALB/c origin (Brittain *et al.*, (1980) *Cancer Res.* 40:179-184), and on B16, a melanoma line derived from C57 B6 mice. Both the CT26 line and B16 are stably transformed to
 30 express human and murine 5T4. Mice are injected I.V. (to induce lung nodules, CT26) or subcutaneously (CT26 and B16) to make single mass subcutaneous tumours.

Groups of 7 BALB/c mice were inoculated three times IV or IM with 1×10^7 pfu of MVA-h5T4 (here the 5T4 antigen is called OBA1) construct on days 0, 21 and 42. Mice were then challenged IV with 5×10^5 tumour cells that were stably transfected with human 5T4. 14 days after challenge mouse lungs were removed and lung nodules counted.

Results 3

The results shown in Figures 3a and 3b demonstrate that mice vaccinated with MVA-h5T4 showed anti-tumour activity when challenged with the syngeneic tumour line CT26 expressing the h5T4 protein. Mann-Whitney statistical analysis show that protection after vaccination with MVA-h5T4 is significant, compared to vaccination with MVA-LacZ or PBS ($p < 0.05$).

Example 4

Groups of 5 C57 BL 6 mice were inoculated twice at a three week interval with 10^7 pfu of MVA-h5T4 (IV or IM) or MVA-Lac Z (IV). Mice were challenged with 5×10^5 B16-h5T4 cells. Tumours sizes were recorded at 2 day intervals post tumour challenge. Individual tumour areas are given.

Groups of 5 and 7 C57 BL 6 mice were inoculated twice at a three week interval with 10^7 pfu of MVA-m5T4 (IV or IM) or MVA-Lac Z (IV). Mice were challenged with 5×10^5 B16-m5T4 cells. Tumours sizes were recorded at 2 day intervals post tumour challenge. Individual tumour areas are given.

Results 4

Figure 4a shows that vaccination with MVA-h5T4 clearly has an anti-tumour effect when mice are challenged with B16-h5T4. Mann-Whitney statistical analysis of data in Figure 4a demonstrates that tumour retardation after vaccination with MVA-h5T4 is significant, compared to vaccination with MVA-LacZ ($p < 0.05$).

Figures 4b and 4c show that vaccination with MVA-m5T4 clearly has an anti-tumour effect when mice are challenged with B16-m5T4. Mann-Whitney statistical analysis of data in Figure 4c demonstrates that tumour retardation after vaccination with MVA-m5T4 is significant, compared to vaccination with MVA-LacZ ($p < 0.05$).

5

Example 5

Female BALB/c mice were injected IV with 5×10^5 CT26-h5T4 cells. After 3 days macro lung tumours establish. Mice were treated on day 3 and 10 post tumour inoculation with 10^7 pfu of MVA-Lac Z, MVA-h5T4 (groups of 10 mice) or PBS (group of 5 mice). Lungs are stained and tumours counted 14 days post tumour inoculation.

10

Results 5

15

It is clear from Figure 5 that treatment with MVA-h5T4 has a significant therapeutic effect on established CT26 lung nodules expressing h5T4. Statistical analysis (Mann-Whitney) shows that therapy with MVA-h5T4 is significant when compared to MVA-Lac Z or PBS ($p < 0.05$).

20

Example 6

CT26-m5T4 and B16-m5T4 Self Antigen Model

C57 BL6 mice are inoculated I.V. twice at a three week interval with 1×10^7 pfu of either MVA-LacZ ($n=3$) or MVA-m5T4 ($n=6$). Three weeks after the last vaccination mice are challenged S.C. with 5×10^5 B16 expressing m5T4. Development of sub cutaneous (S.C.) tumours is monitored.

25

Figure 6 shows that mice that are vaccinated with MVA-m5T4 develop tumours at a slower rate than those that receive the control vaccine. Additionally, the tumours in the m5T4 vaccinated mice are on average 5 fold smaller in volume (10 fold smaller by

30

day 13) compared to those mice that receive the MVA-LacZ treatment. This protective property can be paralleled by the m5T4 antibody response induced in these mice.

5 Example 7

BALB/c mice are inoculated I.V. twice at a three week interval with 1×10^7 pfu of either MVA-LacZ (n=5) or MVA-m5T4 (n=6). Three weeks after the last vaccination mice are challenged with 5×10^5 CT26 expressing m5T4. 12 days after challenge mouse
 10 lungs are removed and tumour nodules counted in a blinded manner. Results 7
 Figure 7 shows that mice that were vaccinated with MVA-m5T4 had a lower tumour burden than those mice that received the MVA-LacZ treatment. This protective property can be paralleled by the m5T4 antibody response induced in these mice.

15 Example 8

Induction of 5T4 Antibody Responses in C57 and BALB/c Mice

Mice are vaccinated as above with MVA-m5T4 and MVA-h5T4, and bled 10 days
 20 after each vaccination.

Results 8

MVA-m5T4 is able to overcome tolerance in both BALB/c and C57 Bl 6 mice, after
 25 two inoculations. Additionally, mice primed with DNA followed by MVA do not show signs of an antibody response to m5T4.

It is therefore shown, in two murine tumour models, that vaccination with MVA
 expressing murine 5T4 has protective properties against syngeneic tumour cells
 30 expressing m5T4. These anti-tumour properties can be paralleled to the anti-m5T4 immune response (measured by ELISA). Additionally, it is shown that the induction of such an immune response does not induce auto immune toxicity in these animals.

Example 9

Immune Response to 5T4 Using Primer Boost Vaccination

5

In order to evaluate the efficacy of MVA and naked DNA vectors to induce immunity to 5T4 in BALB/c and C57 BL6 mice, mice are inoculated using successive priming and boosting with both naked DNA and MVA vectors encoding mouse and human 5T4. In more detail, mice are inoculated with 1×10^7 pfu of MVA-5T4 i.v., 50 g pCl-h5T4 (25 g/hind leg) or 25 g pCl-m5T4 (12.5 g/hind leg) on day 0, with a second (booster) inoculation on day 21 with MVA-5T4. On day 29 the mice are bled and antibody titres determined by ELISA.

10

Results 9

15

The following titres (defined as that dilution giving an OD above twice the background OD of MVA-LacZ) as shown in Tables 2 and 3 are observed:

20

Table 2: Murine 5T4 Antibody Titres from Mice Inoculated with MVA and DNA vectors expressing human and murine 5T4

BALB/c	1:10 000	<1: 1 000	<1: 1 000	<1: 1 000
C57-BL6	1: 16 000	<1: 1 000	<1: 1 000	<1: 1 000

25

Table 3: Human 5T4 Antibody Titres from Mice Inoculated with MVA and DNA vectors expressing human and murine 5T4.

BALB/c	1: 5 000	<1: 1 000	>1: 32 000	1: 32 000
G57 BL6	1: 4 000	<1: 1 000	>1: 32 000	>1: 32 000

The results show that use of either DNA:MVA prime:boost with a heterologous 5T4 antigen (h5T4) or an MVA:MVA prime:boost with heterologous or homologous 5T4 (H5T4 or m5T4) is effective in raising a high titre of antibodies.

5

Example 10

Modified forms of 5T4 for cancer immunotherapy

10 ~~It is possible to modify human 5T4 to enhance its immunogenicity and thus induce more efficacious immunotherapy responses. In order to do this, identification of HLA CTL epitopes and modification of such epitopes to improve binding to the HLA molecule and thus induce more efficient CTL induction is performed using the programme "Peptide Binding Predictions" devised by K. Parker at the National~~
15 ~~institutes of health (http://www.bimas.dcrn.nih.gov/cgi-bin/molbio/ken_parker_comboform) (see Parker, K. C *et al.* 1994.J.Immunol. 152:163). The following results are obtained for human (Table 4) and murine (Table 5) 5T4 9mers:~~

20

Table 4: Human 5T4 9mers binding to HLA A 0201

Rank	Start	Sequence	Dissociation half time
1	97	FLTGNQLAV	319.939
2	364	ALIGAIFLL	284.974
3	351	SLQTSYVFL	176.240
4	368	AIFLLVLYL	137.482
5	283	GLPHIRVFL	117.493
6	358	FLGIVLALI	110.379

7	81	NLTEVPTDL	87.586
8	95	NLFLTGNQL	79.041
9	222	FLYLPRDVL	63.174
10	373	VLYLNRKGI	56.754
11	365	LIGAIFLLV	30.890
12	290	FLDNNPWVC	28.109
13	301	HMADMVTWL	27.207

Table 5: Murine 5T4 binding to human HLA A 0201

Rank	Start	Sequence	Dissociation
1	307	YMADMVAWL	3680.892
2	81	NLLEVPADL	324.068
3	97	FLTGNQMTV	319.939
4	370	ALIGAIFLL	284.974
5	228	FLFLPRDLL	178.158
6	357	SLQTSYVFL	176.240
7	374	AIFLLVLYL	137.482
8	289	GLAHVKVFL	117.493
9	364	FLGIVLALI	110.379
10	379	VLYLNRKG	56.754

5 Example 11

Mutation of H5T4 to improve binding of HLA A0201

Sub. B2
 10 The above data derived from the Parker Peptide Binding Predictions Programme indicates that mutation of the human AA sequence starting at position 301 from YMADMVAWL when changed to HMADMVTWL leads to a 9mer with a 10 fold increase in half time of dissociation to HLA A0201. This increased binding affinity

5

10 Example 12

15

20

Groups of 5 BALB/c and C57 BL6 mice were inoculated IV with 10^7 pfu of
25 recombinant vaccinia virus MVA expressing human (MVA-h5T4) and murine 5T4
(MVA-m5T4). As a negative control mice were inoculated with MVA expressing
E. Coli LacZ (MVA-LacZ) or PBS.

Female BALB/c mice were inoculated a total of four times over an 14-month period.
30 C57 BL6 mice were inoculated 3 times over an 14 month period. Blood samples were
taken after inoculation and evaluated for 5T4 specific antibody by ELISA.

Results 12a

Antibody response: After 2 inoculations mice inoculated with MVA-m5T4 and MVA-h5T4 had high levels of anti-m5T4 and anti-h5T4 respectively, in their serum (see Table 2 and Table 3).

Toxicity: Mice were observed for signs of ill health on a daily basis. At no time during the past 14 months did the physical appearance of animals inoculated with MVA-m5T4 or MVA-h5T4 differ from those animals inoculated with MVA-Lac Z or PBS. Two reports on the health of the animals, prepared by a qualified veterinarian, were prepared that states all animals appear healthy.

Summary

Groups of BALB/c and C57 BL6 mice were inoculated up to four times over an 14 month period with MVA expressing m5T4 (MVA-m5T4) or MVA expressing h5T4 (MVA-h5T4) antigens and checked for signs of toxicity. Though mice were shown to have high titres of antibodies to m5T4 there were no signs of ill health over the 14-month period. A qualified veterinarian has assessed animals and found them to show no signs of ill health, indicating an absence of auto immune toxicity.

Thus, inoculation of BALB/c or C57 BL6 mice with MVA-h5T4 or MVA-m5T4 induces an antibody response to h5T4 and m5T4 respectively. Such a response has no detrimental effect on the health of the mice.

Example 12b

Effects of 5T4 Auto-immunity on Fertility

5T4 is found on human and murine placenta. Accordingly, is possible that an anti mouse 5T4 immune response may prevent mice becoming pregnant or effect the health of the foetus. We carried out extensive studies to address these issues. The purpose of

this study was to assess the effects of an immune response in both BALB/c and C57 BL6 female mice to m5T4 on pregnancy.

Experimental Design

5

Groups of 5 female BALB/c and C57 BL6 mice were inoculated on three occasions IV with 10^7 pfu of MVA-m5T4, MVA-LacZ or PBS. At specific times following the final inoculations (day 10, 30 and 60) mice were mated and evaluated for their ability to become pregnant and give birth to healthy pups.

10

Results 12b

(i) C57 BL6 Mouse Study

15 **Table 6: 10 DAY Study**

	MVA-m5T4 X 3	MVA-LacZ X 3	PBS
No. of pregnancies	4/5 = 80 %	4/5 = 80 %	3/5 = 60 %
No. Live births	24	30	18
Average litter size	6	7.5	6

Table 7: 30 DAY Study

	MVA-m5T4 X 3
No. of pregnancies	4/5 = 80 %
No. Live births	22
Average litter size	5.5

Table 8: 60 DAY Study

	MVA-m5T4 X 3
No. of pregnancies	5/5 = 100 %
No. Live births	33
Average litter size	6.6

20

(ii) BALB-c Mouse Study**Table 9: 10 DAY Study**

	MVA-m5T4 X 3	MVA-LacZ X 3	PBS
No. of pregnancies	5/5 = 100 %	4/5 = 80 %	4/5 = 80 %
No. Live births	30	23	22
Average litter size	6	5.75	5.5
No. surviving to weaning	24 = 80.0 %	19 = 82.6 %	22 = 100 %
Ratio F:M	13:11	14:5	11:11
Average weights	10.4 g	11.2 g	10.1 g

Table 10: 45 DAY Study

	MVA-m5T4 X 3
No. of pregnancies	4/4 = 100 %
No. Live births	24
Average litter size	6
No. surviving to weaning	24 = 100.0 %
Ratio F:M	15:9
Average weights	11.4 g

5

Summary

10

BALB/c and C57 BL6 mice were injected with an MVA recombinant virus expressing m5T4 and an anti-m5T4 antibody response was induced. These mice were mated and were shown to get pregnant and give birth to pups at the same rate as mice that were vaccinated with a control virus (see tables 6-10). Additionally, there was no effect on the health of the pups to weaning. Thus, inoculation with MVA-m5T4 and induction

15

of m5T4 antibody response does not have a detrimental effect on (i) Fertility of BALB/c and C57 BL6 mice; (ii) Number of live births and (iii) Weight and survival of pups to weaning.

5 **Example 12c**

Distribution of 5T4 in Normal Human Tissues

The purpose of this study was to carry out an independent evaluation of 5T4 distribution in normal human tissues. Some past studies have suggested that normal tissue expressing 5T4 could potentially be a target for an immune response induced against this tumour antigen even though the level of 5T4 expression in normal tissue associated with small vessels was found to be over 1000-fold lower than that associated with placenta. It was not clear whether the staining was specific or reflected some cross reactivity by the MAb.

15

Experimental Design

Slide preparations of thirty two different tissue types from 3 different donors were evaluated by a qualified pathologist under GLP conditions. Cryosections of each tissue sample were stained with three different concentrations of Mab specific for 5T4.

20

Results 12c

The summary Table 11 indicates that tissue sections from all essential organs including: brain, CNS, liver and kidney were negative for 5T4 expression. Some weak staining in one or more of the donor tissues was evident in several non-essential tissues. It should be noted that some of the positive staining was observed in tissues derived from individuals that had died from cancer.

25

Although past studies using immunohistochemical analysis have revealed that some 5T4 staining was observed in "some small vessels" of some non-cancerous organs which "appeared to be weakly staining" and include; Kidney (glomeruli), bladder

30

(epithelium), small intestine (villous epithelium), uterus (endometrial glands), cervix (endocervical glands) and skin (basal epidermis), this independent study shows that 5T4 is not expressed on cells of essential organs, furthermore, expression in some normal tissues is at a low level and is sporadic as it is rarely seen in all three donor samples. Thus, this data indicates that 5T4 tissue expression is more stringently restricted compared to some other tumour antigens that have been employed in tumour immunotherapy trials. Therefore these findings reiterate the view that 5T4 is an excellent candidate antigen for cancer immunotherapy.

[illegible]

Table 11: Summary of 5T4 Tissue Distribution

	Tissue	¹ Donors			Staining	Distribution
		1	2	3		
5	Adrenal	-	-	-		
	Bladder	+/++	-	-	Urothelium only stained	(Nt and 1:8)
	Blood Cells					
	Bone Marrow					
10	Breast	-	-	-		
	Brain Cerebellum	-	-	-		
	Brain Cortex	-	-	-		
	Colon	-	-	-		
15	Endothelium	-	+	-	Variable staining in endothelium	
	Fallopian Tube	+	-	-	Variable staining of epithelium	
	Heart	-	-	-		
	Kidney	-	-	-		
20	Liver	-	-	-		
	Uterus Cervix	-	-	-		
	² Uterus Endometrium	-	-	+	Epithelial cells	positive
	Lung	-	-	-		
25	Lymph Node	-	-	-		
	Ovary	+	-	-	Mesothelial and epithelial staining	
	Pancreas	-	-	-		
	Parathyroid	-	-	-		
30	Pituitary	+	-	+	Individual cells	staining
	Placenta	+++	+++	+++	All staining on trophoblasts surface	
	Prostate	-	-	-		
	Skin	-	-	-		
35	Spinal Cord	-	-	-		
	Spleen	-	-	-		
	Striated Muscle	-	-	-		
	Testis	-	-	-		
35	Thymus					
	Thyroid	-	-	-		
	² Ureter	+	+	+	Urothelium only stained	(Nt and 1:8)
	Gastric Antrum	-	-	-		

	Gastric Body	-	-	-	
	Ileum	-	-	-	
	² Duodenum	++/+	-	+	Muscularis mucosa only stained
	Eye Cornea	-	-	-	
5	Eye Lens	-	-	-	
	Eye Retina	-	-	-	

¹5T4 staining intensity was defined by visual analysis relative to placenta trophoblasts (+++) and negative control staining (-).

² Tissues derived from cancer patients

10

Example 13

ScFv Fusion Protein *In vivo* Anti Tumour Efficacy Data

- 15 The purpose of the study was to test the efficacy of a series of single chain antibody fusion proteins.

Experimental Design

- 20 CT26 cells expressing human 5T4 (CT26-h5T4) and CT26-neo

Cells were pre-incubated with:

PBS, LscFv-1, LscFv-2, B7-scFv, ScFv-Ig

- 25 LscFv-1 and 2 were expressed in a BHK cell line. LscFv-1 was purified via its Histidine tag on a Nickel column and scFv-2 was purified using a filtration system. B7-scFv was purified from a BHK line via a His tag and scFv-Ig was purified via a filtration column. The concentration of each scFv used in the experiment was defined as the amount of protein required to saturate binding of CT26-h5T4 cells in a FACS
30 assay.

CT26-h5T4 and CT26-neo cells were pre-incubated with saturating amounts of each scFv and incubated for 1 hour. After washing cells 5×10^5 cells were injected subcutaneously into the flanks of syngeneic BALB/c mice.

- 5 Tumour measurements were taken every two days and the volume calculated.

Results 13

- 10 Figure 8: CT26-neo

There is not a significant difference between the groups except in the case of LscFv-1, for which appears to be 3-fold reduction in tumour size compared to the PBS control 36 days after tumour inoculation.

15

- Figure 9: CT26-h5T4

Tumours treated with all the scFv constructs had a significant effect on tumour growth. 4 of the 5 mice treated with scFv-1 were tumour free on day 36. On day 36 scFv-1 treated tumours were >60 fold smaller than tumours treated with PBS.

20

When a similar experiment was carried out using a mouse melanoma line (B16) engineered to express h5T4 there was no anti-tumour effect.

- 25 *Summary*

In summary there appears to be no benefit of fusing B7 or IgG to the 5T4 specific scFv in the CT26 and B16 murine models. The scFv alone may be more efficacious due to its higher binding affinity (as shown in BIAcore compared to B7-scFv). Therefore this data indicates that the scFv alone has a significant effect on tumour retardation and immune enhancing molecules fused to the scFv may not be required to show an effect on tumour retardation in the 5T4 model.

*Insert
seq. list
per p. #7/Bhuc* →